

The Ando-Hopkins-Rezk orientation is surjective

S. K. Devalapurkar

ABSTRACT. We show that the map $\pi_*\mathrm{MString} \rightarrow \pi_*\mathrm{tmf}$ induced by the Ando-Hopkins-Rezk orientation is surjective. This proves an unpublished claim of Hopkins and Mahowald. We do so by constructing an \mathbf{E}_1 -ring B and a map $B \rightarrow \mathrm{MString}$ such that the composite $B \rightarrow \mathrm{MString} \rightarrow \mathrm{tmf}$ is surjective on homotopy. Applications to differential topology, and in particular to Hirzebruch's prize question, are discussed.

1. Introduction

The goal of this paper is to show the following result due to Hopkins and Mahowald.

THEOREM 1.1 (Hopkins-Mahowald). *The map $\pi_*\mathrm{MString} \rightarrow \pi_*\mathrm{tmf}$ induced by the Ando-Hopkins-Rezk orientation is surjective.*

This integral result was originally stated as [Hop02, Theorem 6.25], but, to the best of our knowledge, no proof has appeared in the literature. In [HM02], Hopkins and Mahowald give a proof sketch of Theorem 1.1 for elements of $\pi_*\mathrm{tmf}$ of Adams-Novikov filtration 0.

The analogue of Theorem 1.1 for bo (namely, the statement that the map $\pi_*\mathrm{MSpin} \rightarrow \pi_*\mathrm{bo}$ induced by the Atiyah-Bott-Shapiro orientation is surjective) is classical [Mil63]. In Section 2, we present (as a warmup) a proof of this surjectivity result for bo via a technique which generalizes to prove Theorem 1.1. We construct an \mathbf{E}_1 -ring A with an \mathbf{E}_1 -map $A \rightarrow \mathrm{MSpin}$. The \mathbf{E}_1 -ring A is a particular \mathbf{E}_1 -Thom spectrum whose mod 2 homology is given by the polynomial subalgebra $\mathbf{F}_2[\zeta_1^4]$ of the mod 2 dual Steenrod algebra. The Atiyah-Bott-Shapiro orientation $\mathrm{MSpin} \rightarrow \mathrm{bo}$ is an \mathbf{E}_∞ -map, and so the composite $A \rightarrow \mathrm{MSpin} \rightarrow \mathrm{bo}$ is an \mathbf{E}_1 -map. We then prove that the map $\pi_*A \rightarrow \pi_*\mathrm{bo}$ is surjective; this is stronger than the Atiyah-Bott-Shapiro orientation being surjective on homotopy.

The argument to prove Theorem 1.1 follows the same outline: we construct (in Section 3) an \mathbf{E}_1 -ring B with an \mathbf{E}_1 -map $B \rightarrow \mathrm{MString}$. This \mathbf{E}_1 -ring B is the height 2 analogue of the \mathbf{E}_1 -ring A ; this motivated the naming of B . We define B as a particular \mathbf{E}_1 -Thom spectrum whose mod 2 homology is given by the polynomial subalgebra $\mathbf{F}_2[\zeta_1^8, \zeta_2^4]$ of the mod 2 dual Steenrod algebra. The Ando-Hopkins-Rezk orientation [AHR10] $\mathrm{MString} \rightarrow \mathrm{tmf}$ is an \mathbf{E}_∞ -map, and so the composite $B \rightarrow \mathrm{MString} \rightarrow \mathrm{tmf}$ is an \mathbf{E}_1 -map. We then prove the following stronger statement:

THEOREM 1.2. *The map $\pi_* B \rightarrow \pi_* \mathrm{tmf}$ is surjective.*

The map $B \rightarrow \mathrm{tmf}$ factors through $\mathrm{MString}$, so Theorem 1.1 follows. In Section 4, we prove Theorem 1.2 after localizing at 3 (as Theorem 4.1). In Section 5, we prove Theorem 1.2 after localizing at 2 (as Theorem 5.1); together, these yield Theorem 1.1 by Corollary 3.6. Finally, in Section 6, we study some applications of Theorem 1.1. In particular, we discuss Hirzebruch’s prize question [HBJ92, Page 86] along the lines of [Hop02, Corollary 6.26]. We also prove a conjecture of Baker’s from [Bak17].

The surjectivity of the Atiyah-Bott-Shapiro orientation $\mathrm{MSpin} \rightarrow \mathrm{bo}$ was considerably strengthened by Anderson, Brown, and Peterson in [ABP67]: they showed that the Atiyah-Bott-Shapiro orientation $\mathrm{MSpin} \rightarrow \mathrm{bo}$ in fact admits a spectrum-level splitting. It is a folklore conjecture that the same is true of the Ando-Hopkins-Rezk orientation $\mathrm{MString} \rightarrow \mathrm{tmf}$, and there have been multiple investigations in this direction (see, for instance, [Lau04, LS19]). In forthcoming work [Dev20], we study in detail the relationship between B and tmf (as well as A and bo). We show that old conjectures of Cohen, Moore, Neisendorfer, Gray, and Mahowald in unstable homotopy theory related to the Cohen-Moore-Neisendorfer theorem, coupled with a conjecture about the centrality of a certain element of $\sigma_2 \in \pi_{13}(B)$ (resp. $\sigma_1 \in \pi_5(A)$), implies that the Ando-Hopkins-Rezk orientation (resp. the Atiyah-Bott-Shapiro orientation) admits a spectrum level splitting. This provides another proof of Theorem 1.1, assuming the truth of these conjectures.

Acknowledgements. I’m extremely grateful to Mark Behrens and Peter May for agreeing to work with me this summer and for being fantastic advisors, as well as for arranging my stay at UChicago. I’d like to also thank Stephan Stolz for useful conversations when I visited Notre Dame. I’m also grateful to Andrew Baker, Hood Chatham, Jeremy Hahn, Eleanor McSpirit, and Zhouli Xu for clarifying discussions. Thanks also to Andrew Baker, Peter May, Haynes Miller, Zhouli Xu, and in particular Jeremy Hahn for providing many helpful comments and correcting mistakes, and to Andrew Senger for pointing out the reference [HM02] after this paper was written. The influence of the ideas of Mike Hopkins and Mark Mahowald on this paper cannot be overstated; Mike told me that the argument presented here is very similar to Mahowald’s and his original proof of Theorem 1.2. I am very grateful to Mike Hopkins for his support and for the influence of his ideas.

2. Warmup: surjectivity of the Atiyah-Bott-Shapiro orientation

The goal of this section is to provide a proof of the following classical theorem using techniques which generalize to prove Theorem 1.1.

THEOREM 2.1. *The map $\pi_* \mathrm{MSpin} \rightarrow \pi_* \mathrm{bo}$ induced by the Atiyah-Bott-Shapiro orientation is surjective.*

As mentioned in the introduction, we prove Theorem 2.1 by constructing an \mathbf{E}_1 -ring A with an \mathbf{E}_1 -map $A \rightarrow \mathrm{MSpin}$. Composing with the Atiyah-Bott-Shapiro orientation $\mathrm{MSpin} \rightarrow \mathrm{bo}$ produces an \mathbf{E}_1 -map $A \rightarrow \mathrm{MSpin} \rightarrow \mathrm{bo}$. We then show the following result, which implies Theorem 2.1.

THEOREM 2.2. *The map $\pi_* A \rightarrow \pi_* \mathrm{bo}$ is surjective.*




FIGURE 1. The 15-skeleton of A at the prime 2 shown horizontally, with 0-cell on the left. The element σ_1 is depicted.

REMARK 2.3. Theorem 2.2 is an old result of Mahowald's: it appears, for instance, as [Mah78, Proposition 4.1(c)] and [HM02, Proposition 2.2(3)]. Theorem 2.2 also implies the second part of [Bak18, Proposition 4.10].

The definition of the \mathbf{E}_1 -ring A is as follows.

CONSTRUCTION 2.4. Let $S^4 \rightarrow \mathrm{BSpin}$ be a generator of $\pi_4 \mathrm{BSpin} \cong \mathbf{Z}$. Since BSpin is an infinite loop space, there is an induced map $\Omega S^5 \rightarrow \mathrm{BSpin}$. Let A denote the Thom spectrum of this map. This is an \mathbf{E}_1 -ring with an \mathbf{E}_1 -map $A \rightarrow \mathrm{MSpin}$. Its 15-skeleton is shown in Figure 1.

REMARK 2.5. The image of a generator of $\pi_4 \mathrm{BSpin}$ under the J-homomorphism $\mathrm{BSpin} \rightarrow \mathrm{BGL}_1(\mathbb{S})$ is the Hopf element $\nu \in \pi_4 \mathrm{BGL}_1(\mathbb{S}) \cong \pi_3 \mathbb{S}$. Consequently, A is the Thom spectrum of the map $\Omega S^5 \rightarrow \mathrm{BGL}_1(\mathbb{S})$ which detects ν on the bottom cell S^4 of the source. In particular, the universal property of Thom spectra from [AB19] exhibits A as the \mathbf{E}_1 -quotient $\mathbb{S} // \nu$ of the sphere spectrum by ν .

REMARK 2.6. The spectrum A is ubiquitous in Mahowald's older works [Mah79, Mah81, Mah82] (where it is often denoted X_5), where its relationship to bo via the composite $A \rightarrow \mathrm{MSpin} \rightarrow \mathrm{bo}$ is utilized to great effect.

PROPOSITION 2.7. *The BP_* -algebra $\mathrm{BP}_*(A)$ is isomorphic to a polynomial algebra $\mathrm{BP}_*[y_2]$, where $|y_2| = 4$. There is a map $A_{(p)} \rightarrow \mathrm{BP}$. On BP -homology, the element y_2 maps to t_1^2 mod decomposables at $p = 2$.*

PROOF. The space ΩS^5 has cells only in dimensions divisible by 4, and hence the same is true of A . The Atiyah-Hirzebruch spectral sequence for $\mathrm{BP}_*(A)$ therefore collapses at the E^2 -page, and so $\mathrm{BP}_*(A) \cong \mathrm{BP}_*[y_2]$, as desired. Since $\pi_* \mathrm{BP}$ is concentrated in even degrees, the element ν vanishes in $\pi_3 \mathrm{BP}$. The universal property of A from Remark 2.5 therefore produces an \mathbf{E}_1 -map $A \rightarrow \mathrm{BP}$. The element ν is detected by $[t_1^2]$ in the 2-local Adams-Novikov spectral sequence for the sphere (in fact, a choice of representative in the cobar complex is $t_1^2 + v_1 t_1$), so this yields the final sentence of the proposition. \square

REMARK 2.8. In particular, the map $A \rightarrow \mathrm{bo}$ is an equivalence in dimensions ≤ 4 . Proposition 2.7 implies that $H_*(A; \mathbf{F}_2) \cong \mathbf{F}_2[\zeta_1^4]$; note that this is the Q_0 -Margolis homology of $H_*(\mathrm{bo}; \mathbf{F}_2) \cong \mathbf{F}_2[\zeta_1^4, \zeta_2^2, \zeta_3, \dots]$. This is sharp: $\pi_5 A$ contains a nontrivial element σ_1 which maps to zero in $\pi_* \mathrm{bo}$. This element is specified up to indeterminacy by the relation $\eta\nu = 0$; see Figure 1.

We will momentarily prove Theorem 2.2. Before doing so, we need to introduce one piece of notation.

NOTATION. Let M be a unital spectrum, and suppose $\alpha, \beta \in \pi_* \mathbb{S}$ and $\gamma \in \pi_* M$ are elements such that $\alpha\beta = 0$ and $\beta\gamma = 0$ in M . Following [BM04, Section

7], the Toda bracket $\langle \alpha, \beta, \gamma \rangle$ will denote the coset of elements of $\pi_{|\alpha|+|\beta|+|\gamma|+1}(M)$ determined by the subgroup $\text{indet} = \alpha\pi_{|\beta|+|\gamma|+1}(M) + \pi_{|\alpha|+|\beta|+1}(\mathbb{S})\gamma$. The subgroup indet is the indeterminacy of the bracket $\langle \alpha, \beta, \gamma \rangle$. There is an analogous definition for higher-fold Toda brackets, but we will not elaborate more on this, since we shall not need it. We will use this notation throughout without further comment.

PROOF OF THEOREM 2.2. Because the map $A \rightarrow \text{bo}$ is one of \mathbf{E}_1 -rings, it suffices to lift all the generators of $\pi_*\text{bo}$ to π_*A . We first prove Theorem 2.2 after inverting 2. Since $\pi_*\text{bo}[1/2] \cong \mathbf{Z}[1/2][u^2]$, where u^2 is the square of the Bott element. It follows from Remark 2.8 that the polynomial generator u^2 of $\pi_*\text{bo}[1/2]$ lifts to $\pi_*A[1/2]$.

It remains to prove Theorem 2.2 after 2-localization. Recall that $\pi_*\text{bo}$ is polynomially generated by η in degree 1 (which is spherical), $2v_1^2$ in degree 4, and v_1^4 in degree 8 (subject to some relations). The element $\eta \in \pi_1\text{bo}_{(2)}$ lifts to $\pi_1A_{(2)}$ because it is a spherical element and the unit $\mathbb{S} \rightarrow \text{bo}_{(2)}$ factors through $A_{(2)}$. It remains to lift the other two generators. Remark 2.8 already shows that $2v_1^2$ lifts to $\pi_*A_{(2)}$. Alternatively, recall that there is a sole d_3 -differential $d_3(v_1^2) = \eta^3$ in the Adams-Novikov spectral sequence for bo . This implies that $2v_1^2 \in \langle 8, \nu, 1_{\text{bo}} \rangle$, with indeterminacy 0 (mod 2). Since ν vanishes in π_*A , we find that the bracket $\langle 8, \nu, 1_A \rangle$ is well-defined in $\pi_4A_{(2)}$.

For v_1^4 , recall that $\sigma = 0$ in $\pi_*\text{bo}$, and that $v_1^4 \in \langle 16, \sigma, 1_{\text{bo}} \rangle \subseteq \pi_*\text{bo}$ (see [BM04, Lemma 7.3]), where the indeterminacy in this bracket is 0 (mod 2). Note that $\langle 16, \sigma, 1_{\text{bo}} \rangle \subseteq \langle 4, 4\sigma, 1_{\text{bo}} \rangle$. We now observe that the attaching map of the 8-cell of $A_{(2)}$ is given by $\sigma + \widetilde{2\nu}$, where $\widetilde{2\nu} \in \pi_7(C\nu)$ is an element determined by the relation $2\nu^2 = 0 \in \pi_6(\mathbb{S})$; see, for instance, [Bak18, Lemma 4.7]. In particular, this implies that $4\sigma = 0$ in $\pi_*A_{(2)}$, so the bracket $\langle 4, 4\sigma, 1_A \rangle$ is well-defined.

It follows from the above discussion that $2v_1^2$ and v_1^4 lift to $\pi_*A_{(2)}$ up to indeterminacy (and the indeterminacy in $\pi_*\text{bo}$ is 0 (mod 2)). If $2v_1^2 + 4nv_1^2 = 2(2n+1)v_1^2$ lifts to $\pi_*A_{(2)}$ for some $n \in \mathbf{Z}_{(2)}$, then so does $2v_1^2$ since $2n+1$ is a 2-local unit. Arguing similarly for v_1^4 , it follows that the other two generators of $\pi_*\text{bo}_{(2)}$ lift to $\pi_*A_{(2)}$, as desired. \square

3. Defining B

In this section, we will define the \mathbf{E}_1 -ring B mentioned in the introduction and study some of its elementary properties. It is the height 2 analogue of the spectrum A from Section 2. We define B as a Thom spectrum whose mod 2 homology is given by $\mathbf{F}_2[\zeta_1^8, \zeta_2^4]$; notice that this is the Q_0 -Margolis homology of the mod 2 homology of tmf . The spectrum B appeared under the name \overline{X} in [HM02, Section 10]. We will work *integrally* (i.e., without inverting any primes) unless explicitly mentioned otherwise.

CONSTRUCTION 3.1. There is a fiber sequence

$$S^9 = \text{O}(10)/\text{O}(9) \rightarrow \text{BO}(9) \rightarrow \text{BO}(10).$$

There is an element $f \in \pi_{12}\text{O}(10) \cong \mathbf{Z}/12$, which is sent to $2\nu \in \pi_{12}(S^9) \cong \mathbf{Z}/24$ under the boundary homomorphism in the long exact sequence on homotopy. Define

a space BN as the homotopy pullback

$$\begin{array}{ccccc} S^9 & \longrightarrow & BN & \longrightarrow & S^{13} \\ \parallel & & \downarrow & & \downarrow f \\ S^9 & \longrightarrow & \mathrm{BO}(9) & \longrightarrow & \mathrm{BO}(10). \end{array}$$

Let N be the loop space ΩBN . If $S^9 \rightarrow \mathrm{B}^2\mathrm{String}$ denotes the generator of $\pi_8 \mathrm{BString}$, then the composite $S^{12} \xrightarrow{2\nu} S^9 \rightarrow \mathrm{B}^2\mathrm{String}$ is null. The Atiyah-Hirzebruch-Serre spectral sequence shows that the generator of $\mathrm{bstring}^1(S^9)$ extends to $\mathrm{bstring}^1(BN)$, and so there is a map $BN \rightarrow \mathrm{B}^2\mathrm{String}$. The induced loop map $N \rightarrow \mathrm{BString}$ is given by the map of fiber sequences

$$(1) \quad \begin{array}{ccccc} N & \longrightarrow & \Omega S^{13} & \longrightarrow & S^9 \\ \downarrow & & \downarrow & & \downarrow \\ \mathrm{BString} & \longrightarrow & * & \longrightarrow & \mathrm{B}^2\mathrm{String}. \end{array}$$

The Thom spectrum of the map $N \rightarrow \mathrm{BString}$ is the \mathbf{E}_1 -ring B .

Note that B is defined integrally, and that it admits an \mathbf{E}_1 -map $B \rightarrow \mathrm{MString}$ obtained by Thomifying the map $N \rightarrow \mathrm{BString}$.

PROPOSITION 3.2. *The BP_* -algebra $\mathrm{BP}_*(B)$ is isomorphic to a polynomial algebra $\mathrm{BP}_*[b_4, y_6]$, where $|b_4| = 8$ and $|y_6| = 12$. There is a map $B_{(p)} \rightarrow \mathrm{BP}$. On BP -homology, the elements b_4 and y_6 map to t_1^4 and t_2^2 mod decomposables at $p = 2$, and y_6 maps to t_1^3 mod decomposables at $p = 3$.*

PROOF. There is a fiber sequence

$$(2) \quad \Omega S^9 \rightarrow N \rightarrow \Omega S^{13}.$$

The J-homomorphism $\mathrm{BString} \rightarrow \mathrm{BGL}_1(\mathbb{S})$ gives a map $N \rightarrow \mathrm{BString} \rightarrow \mathrm{BGL}_1(\mathbb{S})$. The composite with the map $\Omega S^9 \rightarrow N$ gives a map $\Omega S^9 \rightarrow \mathrm{BGL}_1(\mathbb{S})$. This is the extension of the map $S^8 \rightarrow \mathrm{BGL}_1(\mathbb{S})$ detecting $\sigma \in \pi_7(\mathbb{S})$ along $S^8 \rightarrow \Omega S^9$. By one of the main theorems of [AB19], we find that the Thom spectrum of the map $\Omega S^9 \rightarrow \mathrm{BGL}_1(\mathbb{S})$ is the \mathbf{E}_1 -quotient $\mathbb{S} // \sigma$ of the sphere spectrum by σ .

The fiber sequence (2) exhibits B as the Thom spectrum of a map $\Omega S^{13} \rightarrow \mathrm{BGL}_1(\mathbb{S} // \sigma)$. The induced map $S^{12} \rightarrow \mathrm{BGL}_1(\mathbb{S} // \sigma)$ detects an element $\tilde{\nu} \in \pi_{11}(\mathbb{S} // \sigma)$. This element may be described as follows. The relation $\sigma\nu = 0$ in $\pi_*\mathbb{S}$ defines a lift of $\nu \in \pi_3(\mathbb{S})$ to π_{11} of the 15-skeleton $C\sigma$ of $\mathbb{S} // \sigma$; this is the element $\tilde{\nu}$. Since BP is concentrated in even degrees, the element σ vanishes in $\pi_*\mathrm{BP}$. Consequently, $\tilde{\nu}$ is well-defined, and it, too vanishes in $\pi_*\mathrm{BP}$. The universal property of Thom spectra from [AB19] then supplies an \mathbf{E}_1 -map $B \rightarrow \mathrm{BP}$.

In particular, the Thom isomorphism says that the BP -homology of B is abstractly isomorphic as an algebra to the BP -homology of N . This may in turn be computed by the Atiyah-Hirzebruch spectral sequence. However, the fiber sequence (2) implies that the homology of N is concentrated in even degrees. Since $\pi_*\mathrm{BP}$ is also concentrated in even degrees, this implies that the Atiyah-Hirzebruch spectral sequence calculating $\mathrm{BP}_*(B)$ collapses, and we find that $\mathrm{BP}_*(B) \cong \mathrm{BP}_*[b_4, y_6]$, as desired.

The map $B \rightarrow \mathrm{BP}$ induces a map $\mathrm{BP}_*(B) \rightarrow \mathrm{BP}_*(\mathrm{BP}) \cong \mathrm{BP}_*[t_1, t_2, \dots]$. The element $\tilde{\nu}$ is detected by $[t_2^2]$ in the 2-local Adams-Novikov spectral sequence

for \mathbb{S}/σ , and by $[t_1^3]$ in the 3-local Adams-Novikov spectral sequence for \mathbb{S}/σ . The element $\sigma \in \pi_7(\mathbb{S})$ is detected by $[t_1^4]$ in the 2-local Adams-Novikov spectral sequence for the sphere. This yields the final sentence of the proposition. \square

REMARK 3.3. Proposition 3.2 implies that the mod 2 homology of B is isomorphic to $\mathbf{F}_2[\zeta_1^8, \zeta_2^4]$; note that this is the Q_0 -Margolis homology of $H_*(\mathrm{tmf}; \mathbf{F}_2) \cong \mathbf{F}_2[\zeta_1^8, \zeta_2^4, \zeta_3^2, \zeta_4, \dots]$.

REMARK 3.4. The composite $B \rightarrow \mathrm{MString} \rightarrow \mathrm{tmf}$ is an \mathbf{E}_1 -ring map (since the first map is an \mathbf{E}_1 -ring map by construction, and the second is an \mathbf{E}_∞ -ring map by [AHR10]), and it is an equivalence in dimensions ≤ 12 . This follows from Proposition 3.2.

PROPOSITION 3.5. *The map $B \rightarrow \mathrm{tmf}$ induces a surjection on homotopy after inverting 6.*

PROOF. By [Bau08, Proposition 4.4], $\pi_*\mathrm{tmf}[1/6]$ is a polynomial generator on two generators c_4 and c_6 , in degrees 8 and 12, respectively. Since the map $B[1/6] \rightarrow \mathrm{tmf}[1/6]$ is an \mathbf{E}_1 -map, the map $\pi_*B[1/6] \rightarrow \pi_*\mathrm{tmf}[1/6]$ is a ring map. It therefore suffices to lift the elements c_4 and c_6 to $\pi_*B[1/6]$. This follows from Remark 3.4. \square

As an immediate consequence, we have:

COROLLARY 3.6. *If the maps $\pi_*B_{(3)} \rightarrow \pi_*\mathrm{tmf}_{(3)}$ and $\pi_*B_{(2)} \rightarrow \pi_*\mathrm{tmf}_{(2)}$ are surjective, then Theorem 1.2 is true.*

REMARK 3.7. In [Dev20], we show that B is in many ways analogous to tmf . For instance, it satisfies an analogue of the 2-local Wood equivalence $\mathrm{tmf}_{(2)} \wedge DA_1 \simeq \mathrm{tmf}_1(3)_{(2)}$ from [Mat16], where DA_1 is a certain 8-cell complex: the spectrum $B_{(2)} \wedge DA_1$ is a summand of Ravenel's Thom spectrum $X(4)_{(2)}$. (More precisely, it is the summand $T(2)$ of $X(4)_{(2)}$ obtained from the Quillen idempotent, as studied in [Rav86, Chapter 6.5].)

4. Theorem 1.2 after localizing at 3

In Corollary 3.6, we reduced Theorem 1.2 to showing that the maps $\pi_*B_{(3)} \rightarrow \pi_*\mathrm{tmf}_{(3)}$ and $\pi_*B_{(2)} \rightarrow \pi_*\mathrm{tmf}_{(2)}$ are surjective. Our goal in this section is to study the 3-local case. We shall prove:

THEOREM 4.1. *The map $\pi_*B_{(3)} \rightarrow \pi_*\mathrm{tmf}_{(3)}$ is surjective on homotopy.*

CONVENTION 4.2. We shall localize at the prime 3 for the remainder of this section.

4.1. The Adams-Novikov spectral sequence for tmf . In this section, we review the Adams-Novikov spectral sequence for tmf at $p = 3$; as mentioned in Convention 4.2, we shall 3-localize everywhere. The following result is well-known, and is proved in [Bau08]:

THEOREM 4.3. *The E_2 -page of the descent spectral sequence (isomorphic to the Adams-Novikov spectral sequence) for tmf is*

$$H^*(M_{\mathrm{ell}}; \omega^{\otimes 2*}) \cong \mathbf{Z}_3[\alpha, \beta, c_4, c_6, \Delta^{\pm 1}]/I,$$

where I is the ideal generated by the relations

$$3\alpha = 3\beta = 0, \alpha^2 = 0, \alpha c_4 = \beta c_4 = \alpha c_6 = \beta c_6 = 0, c_4^3 - c_6^3 = 1728\Delta.$$

Moreover, α and β are in the image of the map of spectral sequences from the Adams-Novikov spectral sequence of the sphere to that of tmf , with preimages α_1 and β_1 .

The differentials are all deduced from Toda's relation $\alpha_1\beta_1^3 = 0$ in $\pi_*\mathbb{S}$. There is a d_5 -differential $d_5(\beta_{3/3}) = \alpha_1\beta_1^3$ (the "Toda differential"), where $\beta_{3/3}$ lives in bidegree $(t-s, s) = (34, 2)$; see, e.g., [Rav86, Theorem 4.4.22]. Under the \mathbf{E}_∞ -ring map $\mathbb{S} \rightarrow \mathrm{tmf}$, this pushes forward to the same differential in the Adams-Novikov spectral sequence for tmf . Then:

LEMMA 4.4. *There is a relation $\beta_{3/3} = \Delta\beta$ in the E_2 -page of the Adams-Novikov spectral sequence for tmf .*

PROOF. We explain how to deduce this from the literature. Multiplication by α is an isomorphism in the Adams-Novikov spectral sequence for both the sphere and tmf in stem 34, so it suffices to check that $\alpha\beta_{3/3} = \Delta\alpha\beta$. The class $\alpha_1\beta_{3/3}$ (resp. $\Delta\alpha\beta$) is a permanent cycle in the Adams-Novikov spectral sequence of the sphere (resp. tmf) by the discussion on [Rav86, Page 137]. It is known (see [DFHH14, Chapter 13, page 12]) that $\Delta\alpha\beta$ detects $\alpha_1\beta_{3/3}$ in homotopy. To conclude that they are the same on the E_2 -page of the Adams-Novikov spectral sequence for tmf , it suffices to note that $\alpha_1\beta_{3/3}$ maps to (a unit multiple of) $\Delta\alpha\beta$, as desired. \square

It follows by naturality that there is a d_5 -differential $d_5(\Delta\beta) = \alpha\beta^3$, which gives (by β -linearity):

PROPOSITION 4.5. *In the Adams-Novikov spectral sequence for tmf , there is a d_5 -differential $d_5(\Delta) = \alpha\beta^2$.*

Since $3\alpha = 0$ in the Adams-Novikov spectral sequence of tmf , we must have $d_5(3\Delta) = 3\alpha\beta^2 = 0$. There are no other possibilities for differentials on 3Δ , so it is a permanent cycle. Proposition 4.5 shows that there is a Toda bracket $3\Delta \in \langle 3, \alpha, \beta^2 \rangle$ in $\pi_*\mathrm{tmf}$. This can be expressed by the claim that 3Δ can be expressed a composite

$$S^{24} \rightarrow \Sigma^{20}C\alpha_1 \xrightarrow{\beta^2} \mathrm{tmf},$$

where the first map is of degree 3 on the top cell.

By Δ -linearity, there is also a d_5 -differential $d_5(\Delta^2) = \alpha\beta^2\Delta$, so $3\Delta^2$ lives in the E_6 -page. There are no further possibilities for differentials, so $3\Delta^2$ lives in $\pi_*\mathrm{tmf}$. Again, this shows that $3\Delta^2 \in \langle 3, \Delta\alpha, \beta^2 \rangle$. Finally, we turn to Δ^3 . We have $d_5(\Delta^3) = 3\Delta^2\alpha\beta^2$, so we find that $\Delta^3 \in \langle 3, \Delta^2\alpha, \beta^2 \rangle$. We collect our conclusions in the following:

COROLLARY 4.6. *The following is true in $\pi_*\mathrm{tmf}$:*

- $3\Delta \in \langle 3, \alpha, \beta^2 \rangle$;
- $3\Delta^2 \in \langle 3, \Delta\alpha, \beta^2 \rangle$
- $\Delta^3 \in \langle 3, \Delta^2\alpha, \beta^2 \rangle$.

REMARK 4.7. The indeterminacy of the above Toda brackets in $\pi_*\mathrm{tmf}_{(3)}$ are $3\mathbf{Z}_{(3)}\{3\Delta\}$, $3\mathbf{Z}_{(3)}\{3\Delta^2\}$, and $3\mathbf{Z}_{(3)}\{\Delta^3\}$, respectively.

4.2. The Adams-Novikov spectral sequence for B . In this section, we analyze the ring map $B \rightarrow \mathrm{tmf}$, and show that the generators of $\pi_*\mathrm{tmf}_{(3)}$ lift to $\pi_*B_{(3)}$. By Corollary 3.6, this implies Theorem 4.1. We begin by showing:

PROPOSITION 4.8. *There is an element in the E_2 -page of the Adams-Novikov spectral sequence for B which lifts the element Δ in the E_2 -page of the Adams-Novikov spectral sequence for tmf .*

PROOF. To prove the proposition, we begin by recalling the definition of a representative for the element Δ in the cobar complex computing the E_2 -page of the Adams-Novikov spectral sequence for tmf . The Hopf algebroid $(\mathrm{BP}_*\mathrm{tmf}, \mathrm{BP}_*\mathrm{BP} \otimes_{\mathrm{BP}_*} \mathrm{BP}_*\mathrm{tmf})$ is isomorphic to the elliptic curve Hopf algebroid (A, Γ) presenting the moduli stack of cubic curves by [Mat16, Corollary 5.3]. Recall from [Bau08, Page 16] (or [Sil86, Section III.1]) that for an elliptic curve in Weierstrass form

$$(3) \quad y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x + a_6,$$

we can define quantities

$$b_2 = a_1^2 + 4a_2, \quad b_4 = 2a_4 + a_1a_3, \quad b_6 = a_3^2 + 4a_6, \quad b_8 = a_1^2a_6 + 4a_2a_6 - a_1a_3a_4 + a_2a_3^2 - a_4^2,$$

which allows us to define elements

$$c_4 = b_2^2 - 24b_4, \quad c_6 = -b_2^3 + 36b_2b_4 - 216b_6.$$

The discriminant is

$$\Delta = -b_2^2b_8 - 8b_4^3 - 27b_6^2 + 9b_2b_4b_6.$$

Now, it is known that upon inverting 2, every elliptic curve in Weierstrass form (3) is isomorphic to one of the form

$$(4) \quad y^2 = x^3 + a_2x^2 + a_4x.$$

It follows that the elliptic curve Hopf algebroid is isomorphic to a Hopf algebroid of the form $(A', \Gamma') = (\mathbf{Z}[1/2][a_2, a_4], A'[r]/(r^3 + a_2r^2 + a_4r))$, where I is some ideal consisting of complicated relations, and where the Hopf algebroid structure can be written down explicitly (as in [Bau08, Section 3]). A straightforward calculation proves that the discriminant is then

$$(5) \quad \Delta = a_2^2b_4^2 - 16b_4^3.$$

Turning to B , recall that $\mathrm{BP}_*B \cong \mathrm{BP}_*[b_4, y_6]$. The map $(\mathrm{BP}_*B, \mathrm{BP}_*\mathrm{BP} \otimes_{\mathrm{BP}_*} \mathrm{BP}_*B) \rightarrow (A', \Gamma')$ of Hopf algebroids induced by the map $B \rightarrow \mathrm{tmf}$ sends b_4 to b_4 and y_6 to a_2b_4 mod decomposables. It follows from Equation (5) that the element Δ already exists in the 0-line of the Adams-Novikov spectral sequence for B . Using Sage to calculate the 3-series of the formal group law of the elliptic curve (4), one finds that v_1 is a_2 up to a 3-adic unit. We conclude that

$$c_4 = 4v_1^2 - 24b_4, \quad c_6 = -4v_1^3 - 144y_6.$$

This completes the proof of Proposition 4.8. \square

By Remark 3.4, the elements $c_4, c_6 \in \pi_*\mathrm{tmf}$ lift to π_*B . The key to lifting the other elements of $\pi_*\mathrm{tmf}$ is the following:

THEOREM 4.9. *There is a differential $d_5(\Delta) = \alpha\beta^2$ in the Adams-Novikov spectral sequence for B . Moreover, $\alpha\beta^2$ vanishes in π_*B , and 3Δ is a permanent cycle.*




FIGURE 2. Cell structure of the 20-skeleton of B ; the bottom cell (in dimension 0) is on the left; straight lines are α_1 , and curved lines correspond to α_2 and α_4 , in order of increasing length.

PROOF. The element $\alpha\beta^2$ is detected in filtration 5 in the Adams-Novikov spectral sequence for the sphere. We first check that there is no class above filtration 5 in stem 23 the Adams-Novikov spectral sequence for B . In Figure 2, we depict the 20-skeleton of B . Now, $\alpha\beta^2$ is the first class in filtration 5 in the Adams-Novikov spectral sequence for the sphere, so there are no classes above filtration 5 in stem 23 in the algebraic Atiyah-Hirzebruch spectral sequence (converging to the Adams-Novikov spectral sequence of B). Consequently, there are no classes above filtration 5 in stem 23 of the Adams-Novikov spectral sequence for B . It follows that $\alpha\beta^2$ must be detected in filtration 5 in the Adams-Novikov spectral sequence for B . Moreover, if the d_5 -differential on Δ exists, then it is the longest one (and hence 3Δ is a permanent cycle).

We now prove the d_5 -differential. We first claim that there is no nonzero target for a d_r -differential on Δ for $2 \leq r \leq 4$. Indeed, such a class must live in bidegree $(t-s, s) = (23, r)$, so we only need to check that there are no classes in that bidegree. Such a class can only possibly come from those permanent cycles in the algebraic Atiyah-Hirzebruch spectral sequence which are supported on stems $23 - 8 = 15$, $23 - 12 = 11$, $23 - 16 = 7$, or $23 - 20 = 3$ of the Adams-Novikov spectral sequence of the sphere. The only classes in these stems are in Adams-Novikov filtration 1, so cannot possibly contribute to a class that lives in bidegree $(t-s, s) = (23, r)$ with $2 \leq r \leq 4$. Therefore, the first possibility for a differential on Δ is the d_5 -differential $d_5(\Delta) = \alpha\beta^2$. The existence of this differential is forced by the same differential in the Adams-Novikov spectral sequence for tmf .

Therefore, $\alpha\beta^2$ vanishes in the E_∞ -page of the ANSS for B ; there may, however, be a multiplicative extension causing $\alpha\beta^2$ to be nonzero in π_*B . But multiplicative extensions have to jump filtration, and we established that there are no classes above filtration 5 in stem 23 of the Adams-Novikov spectral sequence for B . Therefore, $\alpha\beta^2 = 0$ in π_*B , as desired. \square

COROLLARY 4.10. *The elements $3\Delta, 3\Delta^2, \Delta^3 \in \pi_*\text{tmf}$ lift to π_*B .*

PROOF. Theorem 4.9 verifies that 3Δ lifts to π_*B and that the brackets in Corollary 4.6 are well-defined in π_*B . This implies that $3\Delta^2$ and Δ^3 in $\pi_*\text{tmf}$ lift to π_*B up to indeterminacy. Remark 4.7 tells us the indeterminacy of the brackets in Corollary 4.6. If $3\Delta^2 + 3n[3\Delta^2] = 3(3n+1)\Delta^2$ (resp. $\Delta^3 + 3n\Delta^3 = (3n+1)\Delta^3$) lifts for some nonzero $n \in \mathbf{Z}_{(3)}$, then so does $3\Delta^2$ (resp. Δ^3) since $3n+1$ is a 3-local unit. \square

The elements $\alpha, \beta, c_4, c_6, 3\Delta, 3\Delta^2, \Delta^3$, and $b = \langle \beta^2, \alpha, \alpha \rangle$ (no indeterminacy) generate the homotopy of tmf . Moreover, $\alpha\beta^2 = 0$ in π_*B and $\alpha^2 = 0$ in the sphere,

so b admits a lift to π_*B . Therefore, all generators of $\pi_*\mathrm{tmf}$ admit lifts to π_*B ; this yields Theorem 4.1.

5. Theorem 1.2 after localizing at 2

Our goal in this section is to prove:

THEOREM 5.1. *The map $\pi_*B_{(2)} \rightarrow \pi_*\mathrm{tmf}_{(2)}$ is surjective on homotopy.*

Together with Theorem 4.1 and Corollary 3.6, this proves Theorem 1.2.

CONVENTION 5.2. We shall localize at 2 throughout this section, unless explicitly mentioned otherwise.

5.1. The Adams-Novikov spectral sequence for tmf . In this section, we review the Adams-Novikov spectral sequence for tmf at $p = 2$. The following result is well-known, and is proved in [Bau08] (see also [Beh20, Proposition 1.4.9]):

THEOREM 5.3. *The E_2 -page of the descent spectral sequence (isomorphic to the Adams-Novikov spectral sequence) for tmf is*

$$H^*(M_{\mathrm{ell}}; \omega^{2*}) \cong \mathbf{Z}_{(2)}[c_4, c_6, \Delta^{\pm 1}, \eta, a_1^2\eta, \nu, \epsilon, \kappa, \bar{\kappa}]/I,$$

where I is the ideal generated by the relations

$$\begin{aligned} 2\eta, \eta\nu, 4\nu, 2\nu^2, \nu^3 &= \eta\epsilon, \\ 2\epsilon, \nu\epsilon, \epsilon^2, 2a_1^2\eta, \nu a_1^2\eta, \epsilon a_1^2\eta, (a_1^2\eta)^2 &= c_4\eta^2, \\ 2\kappa, \eta^2\kappa, \nu^2\kappa &= 4\bar{\kappa}, \epsilon\kappa, \kappa^2, \kappa a_1^2\eta, \\ \nu c_4, \nu c_6, \epsilon c_4, \epsilon c_6, a_1^2\eta c_4 &= \eta c_6, a_1^2\eta c_6 = \eta c_4^2, \\ \kappa c_4, \kappa c_6, \bar{\kappa} c_4 &= \eta^4\Delta, \bar{\kappa} c_6 = \eta^2(a_1^2\eta)\Delta, 1728\Delta = c_4^3 - c_6^2. \end{aligned}$$

REMARK 5.4. The elements c_4 and $2c_6$ are permanent cycles. There is a map $\mathrm{tmf} \rightarrow \mathrm{tmf}_1(3)$, where the target is complex oriented. The elements c_4 and $2c_6$ are nontrivial in $\pi_*\mathrm{tmf}_1(3)$. In fact, the image of the map $\mathrm{tmf} \rightarrow \mathrm{tmf}_1(3)$ consists of the elements c_4 , $2c_6$, $c_4\Delta^k$, and $2c_6\Delta^k$ for $k \geq 1$, so these elements must be permanent cycles in the Adams-Novikov spectral sequence for tmf .

The ANSS for tmf is essentially determined from Toda's relation $\bar{\kappa}\nu^3 = 0$ in $\pi_{29}\mathbb{S}$. We will explain this statement in the rest of this section. The relation $\bar{\kappa}\nu^3 = 0 \in \pi_{29}\mathbb{S}$ is enforced by the differential $d_5(\beta_{6/2}) = \bar{\kappa}\nu^3$ in the ANSS for the sphere (see [Isa14]). Then:

LEMMA 5.5. *There is a relation $\beta_{6/2} = \Delta\nu^2$ in the E_2 -page of the Adams-Novikov spectral sequence for tmf .*

This gives the differential $d_5(\Delta\nu^2) = \bar{\kappa}\nu^3$ in the ANSS for tmf . By ν -linearity, we have $d_5(\Delta) = \bar{\kappa}\nu$. Since $4\nu = 0$ in the E_2 -page of the ANSS, the class 4Δ survives. The relation $4\nu = \eta^3$ forces a d_7 -differential on 4Δ . In summary:

THEOREM 5.6. *There are differentials $d_5(\Delta) = \bar{\kappa}\nu$ and $d_7(4\Delta) = \bar{\kappa}\eta^3$ in the ANSS for tmf , and $\bar{\kappa}\nu = 0$ in $\pi_*\mathrm{tmf}$.*

In particular, since $2\eta = 0$ in the ANSS, 8Δ survives to the E_8 -page. There are no more differentials, so it is a permanent cycle. Theorem 5.6 then shows that there is a Toda bracket $8\Delta \in \langle 8, \nu, \bar{\kappa} \rangle$ in $\pi_*\mathrm{tmf}$; this bracket is well-defined since

$8\nu = 0$ in $\pi_*\mathbb{S}$. This can be expressed by the claim that 8Δ may be expressed as a composite

$$S^{24} \rightarrow \Sigma^{20}C\nu \xrightarrow{\bar{\kappa}} \mathrm{tmf},$$

where the first map is degree 8 on the top cell. Similarly, $\Delta\eta \in \langle \eta, \nu, \bar{\kappa} \rangle$ in $\pi_*\mathrm{tmf}$; this bracket is well-defined since $\eta\nu = 0$ in $\pi_*\mathbb{S}$. Arguing in the same way, and using the spherical relations $2\nu^2 = 0$, $\epsilon\nu = 0$, we find that:

PROPOSITION 5.7. *The following Toda brackets exist in $\pi_*\mathrm{tmf}$:*

- (a) $8\Delta \in \langle 8, \nu, \bar{\kappa} \rangle$;
- (b) $\Delta\eta = \langle \eta, \nu, \bar{\kappa} \rangle$;
- (c) $2\Delta\nu = \langle 2\nu, \nu, \bar{\kappa} \rangle$;
- (d) $\Delta\epsilon = \langle \epsilon, \nu, \bar{\kappa} \rangle$;
- (e) $\Delta\eta\kappa = \langle \eta\kappa, \nu, \bar{\kappa} \rangle$;
- (f) $\Delta\eta\bar{\kappa} = \langle \eta\bar{\kappa}, \nu, \bar{\kappa} \rangle$.

None of these except the first have any indeterminacy.

To describe the other elements in $\pi_*\mathrm{tmf}$, we adopt a slightly different approach from Section 4.1 — we will not bother writing down all the generators of $\pi_*\mathrm{tmf}$ as Toda brackets of spherical elements unless it is convenient/necessary to do so. This is only to streamline exposition, although one can of course work this out at one's own leisure; see Remark 5.11.

The d_5 -differential on Δ forces a differential $d_5(\Delta^k) = k\Delta^{k-1}\bar{\kappa}\nu$. The d_7 -differential $d_7(\Delta^4) = \Delta^3\bar{\kappa}\eta^3$ now implies that the classes $\{\Delta^{8k}, 2\Delta^{8k+4}, 4\Delta^{4k+2}, 8\Delta^{2k+1}\}$ survive to the $E_8 = E_9$ -page. In fact, these are permanent cycles. A simple induction on k shows:

PROPOSITION 5.8. *Up to units, we have*

- (a) $\Delta^{8k} \in \langle 2, \Delta^{8k-1}\eta^3, \bar{\kappa} \rangle$ with indeterminacy $2\mathbf{Z}_{(2)}\{\Delta^{8k}\}$;
- (b) $2\Delta^{8k+4} \in \langle 2, \Delta^{8k+3}\eta^3, \bar{\kappa} \rangle$ with indeterminacy $2\mathbf{Z}_{(2)}\{2\Delta^{8k+4}\}$;
- (c) $4\Delta^{4k+2} \in \langle 2, 2\Delta^{4k+1}\nu, \bar{\kappa} \rangle$ with indeterminacy $2\mathbf{Z}_{(2)}\{4\Delta^{4k+2}\}$;
- (d) $8\Delta^{2k+1} \in \langle 8, \Delta^{2k}\nu, \bar{\kappa} \rangle$ with indeterminacy $8\mathbf{Z}_{(2)}\{8\Delta^{2k+1}\}$.

We now turn to the other generators of $\pi_*\mathrm{tmf}$, listed in [Beh20, Figure 1.2].

PROPOSITION 5.9. *We have the following Toda brackets in $\pi_*\mathrm{tmf}$, each without any indeterminacy:*

- (a) $\Delta^2\nu = \langle \nu, 2\nu\Delta, \bar{\kappa} \rangle$;
- (b) $\Delta^4\eta = \langle \eta, \Delta^3\eta^3, \bar{\kappa} \rangle$;
- (c) $\Delta^4\nu = \langle \nu, \Delta^3\eta^3, \bar{\kappa} \rangle$;
- (d) $\Delta^4\epsilon = \langle \epsilon, \Delta^3\eta^3, \bar{\kappa} \rangle$;
- (e) $\Delta^4\kappa = \langle \kappa, 4\nu, 3\nu, 2\nu, \nu, \bar{\kappa}^4 \rangle$;
- (f) $2\Delta^5\nu = \langle 2\nu, \Delta^4\nu, \bar{\kappa} \rangle$;
- (g) $\Delta^5\epsilon = \langle \epsilon, \Delta^4\nu, \bar{\kappa} \rangle$;
- (h) $\Delta^6\nu = \langle \nu, 2\Delta^5\nu, \bar{\kappa} \rangle$;

REMARK 5.10. We have excluded those elements which can be derived using the multiplicative structure. All other elements (except for $c_4\Delta^k$ and $2c_6\Delta^k$) can be expressed as products of the elements listed in Propositions 5.7, 5.8, and 5.9. Importantly, the proofs of these propositions *only* use $\bar{\kappa}\nu = 0$ in $\pi_*\mathrm{tmf}$ (via Theorem 5.6) and multiplicative relations in the sphere.

REMARK 5.11. There are a lot of interesting multiplicative extensions, described in [Bau08, Section 8], but we will not need them. Each of these relations can be derived essentially only using the d_5 -differential of Theorem 5.6 and the multiplicative structure in the homotopy of the sphere.

We can recast these extensions from the following perspective. The spectrum $C\nu$ is the Thom spectrum of the Spin-bundle over S^4 determined by the generator of $\pi_4\mathrm{BSpin}$. Since BSpin is an infinite loop space, this bundle extends to one over ΩS^5 , and hence over the intermediate James constructions $J_k(S^4)$ for all $k \geq 1$. Let $J_k(S^4)^\mu$ denote the Thom spectrum of this bundle, so $J_1(S^4)^\mu = C\nu$. Since $\{J_k(S^4)\}$ forms a filtered \mathbf{E}_1 -space, we obtain a map $C\nu^{\wedge k} \rightarrow J_k(S^4)^\mu$. Taking the product of $\bar{\kappa} : \Sigma^{20}C\nu \rightarrow \mathrm{tmf}$ with itself k times defines a map

$$\bar{\kappa}^k : \Sigma^{20k} J_k(S^4)^\mu \rightarrow \Sigma^{20k} J_k(S^4)^\mu \wedge \mathrm{tmf} \rightarrow \mathrm{tmf}.$$

Suppose $x \in \pi_*\mathbb{S}$ lifts to a map $S^{4k+|x|} \rightarrow J_k(S^4)^\mu$ which given by x on the top $(4k\text{-dimensional})$ cell of $J_k(S^4)^\mu$. Then the composite $S^{4k+|x|} \rightarrow J_k(S^4)^\mu \wedge \mathrm{tmf}$ defines an element of the form $x\Delta^k \in \pi_{24k+|x|}\mathrm{tmf}$. For instance, we have:

- (a) $\Delta^2\nu \in \langle \nu, 2\nu, \nu, \bar{\kappa}^2 \rangle$;
- (b) $\Delta^4\eta \in \langle \eta, 4\nu, 3\nu, 2\nu, \nu, \bar{\kappa}^4 \rangle$;
- (c) $\Delta^4\nu \in \langle \nu, 4\nu, 3\nu, 2\nu, \nu, \bar{\kappa}^4 \rangle$;
- (d) $\Delta^4\epsilon \in \langle \epsilon, 4\nu, 3\nu, 2\nu, \nu, \bar{\kappa}^4 \rangle$;
- (e) $\Delta^4\kappa \in \langle \kappa, 4\nu, 3\nu, 2\nu, \nu, \bar{\kappa}^4 \rangle$;
- (f) $2\Delta^5\nu \in \langle 2\nu, 5\nu, 4\nu, 3\nu, 2\nu, \nu, \bar{\kappa}^5 \rangle$;
- (g) $\Delta^5\epsilon \in \langle \epsilon, 5\nu, 4\nu, 3\nu, 2\nu, \nu, \bar{\kappa}^5 \rangle$.

The brackets in (b), (c), and (e) appear in [Bau08, Corollary 8.7]. The others may also be obtained by arguing as Bauer does: they are consequences of the bracket $\bar{\kappa} = \langle \nu, 2\nu, 3\nu, 4\nu, \nu, \eta \rangle = \langle \nu, 2\nu, 3\nu, 4\nu, \eta, \nu \rangle$ in $\pi_*\mathrm{tmf}$ (no indeterminacy), stated as [Bau08, Lemma 8.6].

REMARK 5.12. Mark Behrens pointed out to us that Mahowald expected $\bar{\kappa}^7 = 0$ in $\pi_*\mathbb{S}_{(2)}$ (it is known that $\bar{\kappa}^6 = 0$ in $\pi_*\mathrm{tmf}_{(2)}$). It would be interesting to know whether this is related to the existence of Δ^8 in $\pi_*\mathrm{tmf}$ via the approach given in Remark 5.11.

Finally, we prove Proposition 5.9.

PROOF OF PROPOSITION 5.9. We prove this case-by-case.

- (a) Since $d_5(\Delta^2) = 2\Delta\bar{\kappa}\nu$ and $2\nu^2 = 0$ in the ANSS for the sphere, we find that $\Delta^2\nu \in \langle \nu, 2\nu\Delta, \bar{\kappa} \rangle$. We provide the argument for indeterminacy in this case, but not for the others since the argument is essentially the same. The indeterminacy lives in $\bar{\kappa}\pi_{31}\mathrm{tmf} + \nu\pi_{48}\mathrm{tmf}$, but $\bar{\kappa}\pi_{31}\mathrm{tmf} \cong \nu\pi_{48}\mathrm{tmf} \cong 0$.
- (b) Since $d_7(\Delta^4) = \Delta^3\bar{\kappa}\eta^3$, we have $d_7(\Delta^4\eta) = \Delta^3\bar{\kappa}\eta^4 = 0$. Therefore, $\Delta^4\eta \in \langle \eta, \Delta^3\eta^3, \bar{\kappa} \rangle$. This bracket is well-defined because $\Delta^3\eta^3 = 4\Delta^3\nu$ exists in $\pi_*\mathrm{tmf}$, $\eta\nu = 0$ in the sphere, and $\bar{\kappa}\eta^3 = 0$ in tmf .
- (c) Similarly, since $d_7(\Delta^4) = \Delta^3\bar{\kappa}\eta^3$, we have $d_7(\Delta^4\nu) = \Delta^3\bar{\kappa}\eta^3\nu = 0$. Therefore $\Delta^4\nu \in \langle \nu, \Delta^3\eta^3, \bar{\kappa} \rangle$. This bracket is well-defined because $\Delta^3\eta^3$ exists in $\pi_*\mathrm{tmf}$, $\eta\nu = 0$ in the sphere, and $\bar{\kappa}\eta^3$ vanishes in tmf .
- (d) Similarly, since $d_7(\Delta^4) = \Delta^3\bar{\kappa}\eta^3$, we have $d_7(\Delta^4\epsilon) = \Delta^3\bar{\kappa}\eta^3\epsilon = 0$, since $2\epsilon = 0$. Therefore, $\Delta^4\epsilon \in \langle \epsilon, \Delta^3\eta^3, \bar{\kappa} \rangle$. This bracket is again well-defined.
- (e) This is in [Bau08, Corollary 8.7], where $\Delta^4\kappa$ is denoted $e[110, 2]$.

- (f) Since $d_5(\Delta^5) = 5\Delta^4\bar{\kappa}\nu$, we have $d_5(2\Delta^5\nu) = 10\Delta^4\bar{\kappa}\nu^2 = 0$, since $2\nu^2 = 0$. It follows that $2\Delta^5\nu \in 5\langle 2\nu, \Delta^4\nu, \bar{\kappa} \rangle$. This is well-defined because $\Delta^4\nu$ lives in $\pi_*\text{tmf}$, $2\nu^2 = 0$ in the sphere, and $\bar{\kappa}\nu = 0$ in tmf .
- (g) Similarly, since $d_5(\Delta^5) = 5\Delta^4\bar{\kappa}\nu$, we have $d_5(\Delta^5\epsilon) = 5\Delta^4\bar{\kappa}\nu\epsilon = 0$, because $\epsilon\nu = 0$. It follows that $\Delta^5\epsilon \in 5\langle \epsilon, \Delta^4\nu, \bar{\kappa} \rangle$, which is well-defined because $\Delta^4\nu$ lives in $\pi_*\text{tmf}$, $\epsilon\nu = 0$ in the sphere, and $\bar{\kappa}\nu = 0$ in tmf .
- (h) Since $d_5(\Delta^6) = 6\Delta^5\bar{\kappa}\nu$, we have $d_5(\Delta^6\nu) = 6\Delta^5\bar{\kappa}\nu^2 = 0$. We therefore have $\Delta^6\nu \in 3\langle \nu, 2\Delta^5\nu, \bar{\kappa} \rangle$. This is well-defined because $2\nu\Delta^5$ lives in $\pi_*\text{tmf}$, $2\nu^2 = 0$ in the sphere, and $\bar{\kappa}\nu = 0$ in tmf .

□

5.2. The Adams-Novikov spectral sequence for B . In this section, we analyze the ring map $B \rightarrow \text{tmf}$, and show that the generators of $\pi_*\text{tmf}_{(2)}$ lift to $\pi_*B_{(2)}$. Again, we will localize at $p = 2$ throughout.

We begin by showing:

PROPOSITION 5.13. *There is an element in the 0-line of the E_2 -page of the ANSS for B which lifts the element Δ in the E_2 -page of the ANSS for tmf .*

PROOF. We begin by recalling a representative for Δ in the cobar complex for tmf at $p = 2$. Recall from Proposition 4.8 that the Hopf algebroid $(\text{BP}_*\text{tmf}, \text{BP}_*\text{BP} \otimes_{\text{BP}_*} \text{BP}_*\text{tmf})$ is isomorphic to the elliptic curve Hopf algebroid (A, Γ) presenting the moduli stack of cubic curves. As in the 3-complete setting (studied in Proposition 4.8), it is known that upon 2-completion, every elliptic curve in Weierstrass form is isomorphic to one of the form

$$y^2 + a_1xy + a_3y = x^3.$$

Consequently (as in the 3-complete setting), the elliptic curve Hopf algebroid is isomorphic to a Hopf algebroid of the form $(A', \Gamma') = (\mathbf{Z}_2[a_1, a_3], A'[s, t]/I)$, where I is some ideal consisting of complicated relations, and where the Hopf algebroid structure can be written down explicitly (as in [Bau08, Section 3]). A straightforward calculation proves that the discriminant is then

$$(6) \quad \Delta = a_1^3a_3^3 - 27a_3^4 = b_4^3 - 27b_6^2.$$

Turning to B , recall that we may identify BP_*B with $\text{BP}_*[b_4, y_6]$. The map $B \rightarrow \text{tmf}$ induces a map $(\text{BP}_*B, \text{BP}_*\text{BP} \otimes_{\text{BP}_*} \text{BP}_*B) \rightarrow (A', \Gamma')$ of Hopf algebroids that sends b_4 to b_4 and y_6 to b_6 mod decomposables. It follows from Equation (6) that the element Δ already exists in the 0-line of the Adams-Novikov spectral sequence for B . This finishes the proof of Proposition 5.13. □

Since the map $B \rightarrow \text{tmf}$ is an equivalence in dimensions ≤ 12 (Corollary 3.6), the elements c_4 and $2c_6$ lift to π_*B . We claim that $c_4\Delta^k$ and $2c_6\Delta^k$ live in π_*B ; to show this, we argue as in Remark 5.4. There is a map $B \rightarrow B \wedge DA_1 \simeq T(2)$ (see also Remark 3.7), and there is a particular complex orientation of $\text{tmf}_1(3)$ exhibiting it as a form of $\text{BP}\langle 2 \rangle$, which sits in a commutative diagram

$$\begin{array}{ccccc} B & \longrightarrow & T(2) & \longrightarrow & \text{BP} \\ \downarrow & & \downarrow & \swarrow & \\ \text{tmf} & \longrightarrow & \text{tmf}_1(3) & & \end{array}$$

There are choices of indecomposables v_1 and v_2 producing an isomorphism $\pi_*\mathrm{tmf}_1(3) \cong \mathbf{Z}_2[v_1, v_2]$ such that c_4 is sent to v_1^4 and Δ is sent to v_2^4 . The map $T(2) \rightarrow \mathrm{tmf}_1(3)$ is surjective on homotopy, since v_1 and v_2 live in $\pi_*T(2)$. Since the elements c_4 , $2c_6$, $c_4\Delta^k$, and $2c_6\Delta^k$ for $k \geq 1$ therefore already live in the homotopy of $T(2)$, we find by the same argument that these elements already live in the homotopy of B .

We next turn to showing that the other elements of $\pi_*\mathrm{tmf}$ lift to π_*B . The following is the 2-local analogue of Theorem 4.9:

THEOREM 5.14. *There are differentials $d_5(\Delta) = \bar{\kappa}\nu$ and $d_7(4\Delta) = \bar{\kappa}\eta^3$ in the ANSS for B . Moreover, $\bar{\kappa}\nu = 0$ in π_*B , and 8Δ is a permanent cycle.*

PROOF. To prove the differentials, first note that the d_7 -differential follows from the d_5 -differential via the spherical relation $4\nu = \eta^3$; it therefore suffices to prove the d_5 -differential. The class $\bar{\kappa}\nu$ lives in bidegree $(23, 5)$ in the ANSS for B , since it lives in that bidegree in the ANSS for both the sphere and for tmf . We claim that if $\bar{\kappa}\nu^2$ vanishes in π_*B , then the d_5 -differential follows. It suffices to establish that $d_5(\Delta\nu) = \bar{\kappa}\nu^2$, since the desired d_5 -differential then follows from ν -linearity. Since $\bar{\kappa}\nu^2$ is the first element of filtration 5 in the ANSS for the sphere which does not come from an η -tower on the α -family elements (and such η -towers are truncated by ANSS d_3 -differentials), there cannot be any differential off it. Moreover, if it is killed on any finite page in the ANSS, then it must in fact be zero in homotopy, since multiplicative extensions have to jump in filtration (and there is nothing of higher filtration). We need to show that $\bar{\kappa}\nu^2$ cannot be the target of a d_r -differential for $2 \leq r \leq 4$; then the claimed d_5 -differential on the E_5 -page is forced by the same differential in the ANSS for tmf . The algebraic Atiyah-Hirzebruch spectral sequence for the ANSS of B implies that the only possibility for a differential is a d_3 ; but the source of any nontrivial d_3 -differential vanishes when mapped to the ANSS for tmf , so no such d_3 -differential can exist.

We now show that $\bar{\kappa}\nu^2$ vanishes in π_*B . For this, we argue as in [HM14, Proposition 8.1]. Namely, [HM14, Lemma 8.2] states that $\bar{\kappa}\nu^2 \in \langle \eta_4\sigma, \eta, 2 \rangle$. Recall that $\eta_4 = h_1h_4$; by [IWX20, Table 21], there is a σ -extension from h_1h_4 to h_4c_0 . There is no indeterminacy in the above Toda bracket, so $\bar{\kappa}\nu^2$ will vanish if we show that h_4c_0 vanishes in $\pi_{23}(B)$. In fact, it vanishes in the E_2 -page of the ANSS for B : since the attaching map of the 8-cell of B is σ , the σ -extension on η_4 implies that h_4c_0 is killed in the algebraic Atiyah-Hirzebruch spectral sequence for the ANSS of B by a d_1 -differential off the ANSS class h_1h_4 supported on the cell in dimension 8. \square

Finally:

PROOF OF THEOREM 5.1. Theorem 5.14 implies that 8Δ lifts to π_*B , and that all the brackets in $\pi_*\mathrm{tmf}$ in Propositions 5.7, 5.8, and 5.9 are well-defined in π_*B . The elements of $\pi_*\mathrm{tmf}$ in those propositions for which the bracket has no indeterminacy therefore lift to π_*B . By Remark 5.10, all that remains is to show that the constant multiples of the powers of Δ which live in $\pi_*\mathrm{tmf}$ in fact lift to π_*B . Theorem 5.14 implies that they lift up to indeterminacy, and this indeterminacy is specified in Proposition 5.8. If $\Delta^{8k} + 2n\Delta^{8k} = (2n+1)\Delta^{8k}$ lifts for some $n \in \mathbf{Z}_{(2)}$, then so does Δ^{8k} since $2n+1$ is a 2-local unit. Similarly, one finds that $2\Delta^{8k+4}$, $4\Delta^{4k+2}$, and $8\Delta^{2k+1}$ also lift to π_*B , as desired. \square

REMARK 5.15. We briefly look at the Adams spectral sequence for B . The Steenrod module structure of the 20-skeleton of B is as in Figure 2; since we are at the prime 2, straight lines are Sq^4 , and curved lines correspond to Sq^8 and Sq^{16} , in order of increasing length. Using this, we can calculate the Adams spectral sequence in small dimensions. The Adams charts below were created with Hood Chatham's Ext calculator, and the Steenrod module file for B in this range can be found at <https://sanathdevalapurkar.github.io/files/input-B-leq-24-prime-2.md>.

The E_2 -page for B in the first few dimensions is shown in Figure 3; there are no classes in higher Adams filtration in stem 23. The red class is $g = \bar{\kappa}$, and the purple lines are d_2 -differentials. The differential on the class in stem 23 already exists in the Adams spectral sequence for the sphere as $d_2(i) = h_0 P d_0$. The other classes in stem 23 except for the one in filtration 9 are permanent cycles, and there is no multiplicative extension causing any of them to be $\bar{\kappa}\nu$ on homotopy.

As shown in Figure 4, there is also a d_3 -differential on the leftmost class $x_{24,1}^{(0)}$ in bidegree $(24, 6)$ (which supports a h_0 -tower) to the class in bidegree $(23, 9)$; the class $h_0 x_{24,1}^{(0)}$ is a permanent cycle in the ASS for B which is sent to 8Δ in the ASS for tmf . The class in bidegree $(25, 5)$ is a permanent cycle in the ASS for B which is sent to $\Delta\eta$ in the ASS for tmf .

REMARK 5.16. We now compare the approach of this paper with that of [HM02], where the \mathbf{E}_1 -ring B was constructed under the name \bar{X} . The special case of our Theorem 1.2 for elements in $\pi_*\mathrm{tmf}$ of ANSS filtration 0 is stated as [HM02, Theorem 11.1], where a proof is only sketched.

First, their Proposition 11.2 is a combination of our Theorem 4.9 and Theorem 5.14. Secondly, their proof proceeds by calculating the mod 2 Adams spectral sequence of B in dimensions ≤ 24 to show that $\bar{\kappa}\nu$ vanishes in the 2-local homotopy of B . Their argument does not seem to resolve potential multiplicative extensions: as Figure 4 shows, there are two possibilities for multiplicative extensions in the Adams spectral sequence which could make $\bar{\kappa}\nu$ nonzero in $\pi_*B_{(2)}$. (Namely, the classes in bidegrees $(23, 6)$ and $(23, 7)$ could represent $\bar{\kappa}\nu$.) Thirdly, Remark 5.11 essentially gives a proof of their Lemma 11.5, which seems to appear without proof.

6. Applications

In this section, we study some applications of Theorem 1.1 and Theorem 1.2.

6.1. A conjecture of Baker's. In [Bak17], Baker constructed a certain collection of \mathbf{E}_∞ -ring spectra Mj_r with \mathbf{E}_∞ -ring maps $Mj_1 \rightarrow \mathbf{HZ}$, $Mj_2 \rightarrow \mathbf{bo}$, and $Mj_3 \rightarrow \mathrm{tmf}$. He conjectured in [Bak17, Conjecture 6.2] that the map $\pi_*Mj_3 \rightarrow \pi_*\mathrm{tmf}$ is surjective. In this section, we show that this conjecture follows from Theorem 1.2.

We begin by recalling the definition of the \mathbf{E}_∞ -ring spectra Mj_r .

DEFINITION 6.1. Let $\mathrm{BO}\langle 2^r \rangle^{[2^{r+1}-1]}$ denote the $(2^{r+1} - 1)$ -skeleton of $\mathrm{BO}\langle 2^r \rangle$. Since $\mathrm{BO}\langle 2^r \rangle$ is an infinite loop space, the skeletal inclusion $\mathrm{BO}\langle 2^r \rangle^{[2^{r+1}-1]} \rightarrow \mathrm{BO}\langle 2^r \rangle$ induces a map $\Omega^\infty \Sigma^\infty \mathrm{BO}\langle 2^r \rangle^{[2^{r+1}-1]} \rightarrow \mathrm{BO}\langle 2^r \rangle$. The Thom spectrum of this map is the \mathbf{E}_∞ -ring Mj_r .

There is an evident \mathbf{E}_∞ -map $Mj_r \rightarrow \mathrm{MO}\langle 2^r \rangle$, which in the case $r = 3$ defines an \mathbf{E}_∞ -map $Mj_3 \rightarrow \mathrm{MString}$. The following result proves the aforementioned conjecture of Baker's as an application of Theorem 1.2:




FIGURE 3. E_2 -page of the Adams spectral sequence for B . The class highlighted in red is $\bar{\kappa}$.




FIGURE 4. E_3 -page of the Adams spectral sequence for B . The class highlighted in red is $\bar{\kappa}$. There are no differentials in this range from the E_4 -page onwards.

PROPOSITION 6.2 ([Bak17, Conjecture 6.2]). *The composite $Mj_3 \rightarrow \text{MString} \rightarrow \text{tmf}$ is surjective on homotopy, where $\text{MString} \rightarrow \text{tmf}$ is the Ando-Hopkins-Rezk orientation.*

PROOF. By Theorem 1.2, it suffices to show that the map $B \rightarrow \text{MString}$ factors through a map $B \rightarrow MJ_3$. Since Mj_3 is the Thom spectrum of a bundle over $\Omega^\infty \Sigma^\infty \text{BString}^{[15]}$, this in turn follows from the existence of a map $N \rightarrow \Omega^\infty \Sigma^\infty \text{BString}^{[15]}$ factoring $N \rightarrow \text{BString}$. Recall that the map $N \rightarrow \text{BString}$ was constructed via the map (1) of fiber sequences. The map $S^9 \rightarrow \text{B}^2\text{String}$ factors as $S^9 \rightarrow \Omega^{\infty-1} \Sigma^\infty \text{BString}^{[15]}$, and so the map of fiber sequences in (1) factors as

$$\begin{array}{ccccc} N & \longrightarrow & \Omega S^{13} & \longrightarrow & S^9 \\ \downarrow \exists & & \downarrow & & \downarrow \\ \Omega^\infty \Sigma^\infty \text{BString}^{[15]} & \longrightarrow & * & \longrightarrow & \Omega^{\infty-1} \Sigma^\infty \text{BString}^{[15]} \\ \downarrow & & \downarrow & & \downarrow \\ \text{BString} & \longrightarrow & * & \longrightarrow & \text{B}^2\text{String}, \end{array}$$

as desired. \square

6.2. Hirzebruch's prize question. Another application of Theorem 1.1 was stated as [Hop02, Corollary 6.26], and provides an answer to Hirzebruch's prize question [HBJ92, Page 86]. See also [HM02].

COROLLARY 6.3. *There exists a 24-dimensional compact smooth string manifold M with $\hat{A}(M) = 1$ and $\hat{A}(M, \tau_M \otimes \mathbf{C}) = 0$.*

PROOF. By the discussion on [HBJ92, Page 86], the conditions on the \hat{A} -genus of M are equivalent to the Witten genus of M being $c_4^3 - 744\Delta = \Delta(j - 744)$, where j is the j -function. Let M_0^8 denote the Kervaire-Milnor almost parallelizable 8-manifold; then, the 8-manifold $-M_0^8 - 224\mathbf{H}P^2$ (whose string cobordism class we will denote by $[N_{c_4}]$, where N_{c_4} is the explicit manifold representative above) admits a string structure by [Lau04, Lemma 15]. The map $\text{tmf} \rightarrow \text{bo}$ sends $c_4 \in \pi_8 \text{tmf}$ to $v_1^4 \in \pi_8 \text{bo}$. By Lemma 6.4, there is a commutative diagram:

$$(7) \quad \begin{array}{ccc} \text{MString} & \longrightarrow & \text{MSpin} \\ \downarrow & & \downarrow \\ \text{tmf} & \longrightarrow & \text{bo}, \end{array}$$

where the left vertical map is the Ando-Hopkins-Rezk orientation and the right vertical map is the Atiyah-Bott-Shapiro orientation. Consequently, the Witten genus of $-M_0^8 - 224\mathbf{H}P^2$ is c_4 .

By Theorem 1.1, the element $24\Delta \in \pi_{24} \text{tmf}$ lifts to a class $[N_\Delta]$ in $\pi_{24} \text{MString}$, where N_Δ is any manifold representative. Since $744\Delta = 31 \cdot 24\Delta$, we conclude that the string cobordism class of the 24-dimensional compact oriented smooth string manifold $N_{c_4}^3 - 31N_\Delta$ has Witten genus $c_4^3 - 744\Delta$, as desired. \square

The proof of Corollary 6.3 utilized the following lemma.

LEMMA 6.4. *The diagram (7) commutes, where the left vertical map is the Ando-Hopkins-Rezk orientation and the right vertical map is the Atiyah-Bott-Shapiro orientation.*

PROOF. We need to show that the composite $MString \rightarrow tmf \rightarrow bo$ comes from the Atiyah-Bott-Shapiro orientation. By [AHR10, Corollary 7.12], it suffices to show that this composite has the same characteristic series as the restriction of the \hat{A} -genus to string manifolds. There is an isomorphism $\pi_* bo \otimes \mathbf{Q} \cong \mathbf{Z}[\beta^2]$, where β^2 lives in degree 4 and is the square of the Bott element. Moreover, $\pi_* tmf \otimes \mathbf{Q}$ is isomorphic to the ring of rational modular forms (of weight given by half the degree in $\pi_* tmf \otimes \mathbf{Q}$) by [Bau08, Proposition 4.4]. The map $\pi_* tmf \otimes \mathbf{Q} \rightarrow \pi_* bo \otimes \mathbf{Q}$ sends a modular form of weight k with q -expansion $f(q) = \sum a_n q^n$ to the element $a_0(\beta^2)^k \in \pi_{2k} bo \otimes \mathbf{Q}$. Consequently, the composite $\pi_* MString \rightarrow \pi_* tmf \otimes \mathbf{Q} \rightarrow \pi_* bo \otimes \mathbf{Q}$ sends a string manifold M to the constant term of the q -expansion of its Witten genus. The lemma will therefore follow if this constant term is the \hat{A} -genus of M , but this follows from the discussion on [HBJ92, Page 84]. \square

REMARK 6.5. The modular form $c_4^3 - 744\Delta$ is $\theta_{\Lambda_{24}} - 24\Delta$, where Λ_{24} is the 24-dimensional Leech lattice and $\theta_{\Lambda_{24}}$ is its theta function.

REMARK 6.6. The original motivation for Hirzebruch's prize question was to relate the geometry of the 24-dimensional string manifold M of Corollary 6.3 to representations of the monster group by constructing an action of the monster group on M . The question of constructing this action remains unresolved.

REMARK 6.7. The discussion on [HBJ92, Page 86] implies that $\hat{A}(N_\Delta) = 0$ and $\hat{A}(N_\Delta, \tau_{N_\Delta} \otimes \mathbf{C}) = 24$. It follows from [Sto92, Theorem A] that N_Δ (which we may assume is simply-connected by surgery) admits a metric with positive scalar curvature. Since the Witten genus of N_Δ is nonzero, Stolz's conjecture in [Sto96] would imply that it does not admit a metric of positive-definite Ricci curvature. We do not know whether Stolz's conjecture holds in this particular case. Note, however, that there are examples of non-simply-connected manifolds which admit positive scalar curvature metrics but no metrics of positive-definite Ricci curvature: as pointed out to us by Stolz, a connected sum of lens spaces of dimension at least 3 gives such a manifold.

Corollary 6.3 may be generalized in the following manner. Recall the following definition from [Ono04, Section 2.3]. Let $j_1(z) = j(z) - 744$, and define $j_n(z)$ for $n \geq 2$ via $nT_n(j_1(z))$, where T_n is the weight zero Hecke operator, acting on $f(z)$ via

$$T_n f(z) = \sum_{d|n, ad=n} \sum_{b=0}^{d-1} f\left(\frac{az+b}{d}\right).$$

By [Ono04, Proposition 2.13], $j_n(z)$ is a monic integral polynomial in $j(z)$ of degree n ; for instance,

$$j_2(z) = j(z)^2 - 1488j(z) + 159768, \quad j_3(z) = j(z)^3 - 2232j(z)^2 + 1069956j(z) - 36866976.$$

The functions $j_n(z)$ for $n \geq 0$ (where $j_0(z) = 1$) form a basis for the complex vector space of weakly holomorphic modular forms of weight 0, and appear in the denominator formula for the monster Lie algebra. They may be defined by Faber polynomials on j . The generalization of Corollary 6.3 is as follows.

PROPOSITION 6.8. *For all $n \geq 0$, there is a $24n$ -dimensional compact smooth string manifold M^{24n} whose Witten genus is $\Delta^n j_n(z)$.*

REMARK 6.9. By arguing as in [HBJ92, Pages 86-87], we find that the twisted \hat{A} -genera of bundles over M^{24n} constructed from the complexified tangent bundle of M are integral linear combinations of dimensions of irreducible representations of the monster group; for instance, $\hat{A}(M^{48}; \text{Sym}^2(\tau_M \otimes \mathbf{C}))$ is the coefficient of q^2 in $\Delta^2 j_2(z)$, which is $2 \times (21296876 + 196883 + 1)$. More generally, $\hat{A}(M^{24n}; \text{Sym}^2(\tau_M \otimes \mathbf{C}))$ is an integral linear combination of the dimensions of the n smallest irreducible representations of the monster group. In light of Hirzebruch's original motivation for his prize question (see Remark 6.6), it seems reasonable to conjecture that the $24n$ -dimensional string manifold M^{24n} admits an action of the monster group by diffeomorphisms.

REMARK 6.10. It would be interesting to know if there is an analogue of Proposition 6.8 for other McKay-Thompson series.

Before providing the proof, we need the following result.

THEOREM 6.11. *A modular form f is in the image of the boundary homomorphism $\pi_* \text{tmf} \rightarrow \text{MF}_*$ in the Adams-Novikov spectral sequence if and only if it is expressible as an integral linear combination of monomials of the form $a_{ijk} c_4^i c_6^j \Delta^k$ with $i, k \geq 0$ and $j = 0, 1$, where*

$$a_{ijk} = \begin{cases} 1 & i > 0, j = 0 \\ 2 & j = 1 \\ 24/\gcd(24, k) & i, j = 0. \end{cases}$$

PROOF. This is [Hop02, Proposition 4.6], proved in [Bau08]. \square

PROOF OF PROPOSITION 6.8. We have

$$\Delta^n j_n(z) = \sum_{0 \leq k \leq n} \alpha_k j(z)^k \Delta^n = \sum_{0 \leq k \leq n} \alpha_k c_4^{3k} \Delta^{n-k},$$

for some integers α_k (where $\alpha_n = 1$). By Theorem 1.1 and Theorem 6.11, it suffices to show that the constant term α_0 of $j_n(z)$ (when expanded as a monic integral polynomial in $j(z)$) is a multiple of $24/\gcd(24, n)$. The j -function vanishes on a primitive third root of unity, so $\alpha_0 = j_n(\omega)$. Its generating function is

$$\sum_{n \geq 0} j_n(\omega) q^n = -\frac{j'(z)}{j(z)} = \frac{c_6}{c_4},$$

where $q = e^{2\pi iz}$ and ω is a primitive third root of unity.

Let $m \geq 1$; we claim that the coefficients $a_{4,m}$ and $a_{6,m}$ of q^m in the q -expansion for c_4 and c_6 (respectively) are divisible by $24/\gcd(24, m)$. Indeed, the expression for their q -expansion shows that $a_{4,n} = -240\sigma_3(n)$ and $a_{6,n} = 504\sigma_5(n)$, and both 240 and 504 are already divisible by 24. Since the coefficient of q^m in $1/c_4$ can be expressed as an integral linear combination of the $a_{4,k}$, it follows that the coefficient of q^m for $m \geq 1$ in c_6/c_4 (which is $j_m(\omega)$) is divisible by 24, and hence by $24/\gcd(24, m)$, as desired. \square

References

- [AB19] O. Antolín-Camarena and T. Barthel. A simple universal property of Thom ring spectra. *J. Topol.*, 12(1):56–78, 2019.
- [ABP67] D. W. Anderson, E. H. Brown, and F. P. Peterson. The structure of the Spin cobordism ring. *Ann. of Math. (2)*, 86:271–298, 1967.
- [AHR10] M. Ando, M. Hopkins, and C. Rezk. Multiplicative orientations of KO -theory and of the spectrum of topological modular forms. <http://www.math.uiuc.edu/~mando/papers/koandtmf.pdf>, May 2010.
- [Bak17] A. Baker. \mathcal{E}_∞ ring spectra and elements of Hopf invariant 1. *Bol. Soc. Mat. Mex. (3)*, 23(1):195–231, 2017.
- [Bak18] A. Baker. Characteristics for \mathcal{E}_∞ ring spectra. In *An alpine bouquet of algebraic topology*, volume 708 of *Contemp. Math.*, pages 1–17. Amer. Math. Soc., Providence, RI, 2018.
- [Bau08] T. Bauer. Computation of the homotopy of the spectrum tmf . In *Groups, homotopy and configuration spaces*, volume 13 of *Geom. Topol. Monogr.*, pages 11–40. Geom. Topol. Publ., Coventry, 2008.
- [Beh20] M. Behrens. Topological modular and automorphic forms. In *Handbook of homotopy theory*, pages 223–265, 2020.
- [BM04] A. Baker and P. May. Minimal atomic complexes. *Topology*, 43(3):645–665, 2004.
- [Dev20] S. Devalapurkar. Higher chromatic Thom spectra via unstable homotopy theory. <https://arxiv.org/abs/2004.08951>, 2020.
- [DFHH14] C. Douglas, J. Francis, A. Henriques, and M. Hill. *Topological Modular Forms*, volume 201 of *Mathematical Surveys and Monographs*. American Mathematical Society, 2014.
- [HBJ92] F. Hirzebruch, T. Berger, and R. Jung. *Manifolds and modular forms*. Aspects of Mathematics, E20. Friedr. Vieweg & Sohn, Braunschweig, 1992. With appendices by Nils-Peter Skoruppa and by Paul Baum.
- [HM02] M. Hopkins and M. Mahowald. The structure of 24 dimensional manifolds having normal bundles which lift to $BO[8]$. In *Recent progress in homotopy theory (Baltimore, MD, 2000)*, volume 293 of *Contemp. Math.*, pages 89–110. Amer. Math. Soc., Providence, RI, 2002.
- [HM14] M. Hopkins and M. Mahowald. From elliptic curves to homotopy theory. In *Topological modular forms*, volume 201 of *Math. Surveys Monogr.*, pages 261–285. Amer. Math. Soc., Providence, RI, 2014.
- [Hop02] M. J. Hopkins. Algebraic topology and modular forms. In *Proceedings of the International Congress of Mathematicians, Vol. I (Beijing, 2002)*, pages 291–317. Higher Ed. Press, Beijing, 2002.
- [Isa14] D. Isaksen. Classical and motivic Adams-Novikov charts. <https://arxiv.org/abs/1408.0248>, 2014.
- [IWX20] D. Isaksen, G. Wang, and Z. Xu. More stable stems. <https://arxiv.org/abs/2001.04511>, 2020.
- [Lau04] G. Laures. $K(1)$ -local topological modular forms. *Invent. Math.*, 157(2):371–403, 2004.
- [LS19] G. Laures and B. Schuster. Towards a splitting of the $K(2)$ -local string bordism spectrum. *Proc. Amer. Math. Soc.*, 147(1):399–410, 2019.
- [Mah78] M. Mahowald. Some homotopy classes generated by η_j . In *Algebraic topology, Aarhus 1978 (Proc. Sympos., Univ. Aarhus, Aarhus, 1978)*, volume 763 of *Lecture Notes in Math.*, pages 23–37, 1978.
- [Mah79] M. Mahowald. Ring spectra which are Thom complexes. *Duke Math. J.*, 46(3):549–559, 1979.
- [Mah81] M. Mahowald. bo -resolutions. *Pacific J. Math.*, 92(2):365–383, 1981.
- [Mah82] M. Mahowald. The image of J in the EHP sequence. *Ann. of Math. (2)*, 116(1):65–112, 1982.
- [Mat16] A. Mathew. The homology of tmf . *Homology Homotopy Appl.*, 18(2):1–20, 2016.
- [Mil63] J. Milnor. Spin structures on manifolds. *Enseignement Math. (2)*, 9:198–203, 1963.
- [Ono04] K. Ono. *The web of modularity: arithmetic of the coefficients of modular forms and q -series*, volume 102 of *CBMS Regional Conference Series in Mathematics*. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 2004.

- [Rav86] D. Ravenel. *Complex cobordism and stable homotopy groups of spheres*. Academic Press, 1986.
- [Sil86] J. Silverman. *The arithmetic of elliptic curves*, volume 106 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1986.
- [Sto92] S. Stolz. Simply connected manifolds of positive scalar curvature. *Ann. of Math. (2)*, 136(3):511–540, 1992.
- [Sto96] S. Stolz. A conjecture concerning positive Ricci curvature and the Witten genus. *Math. Ann.*, 304(4):785–800, 1996.

Email address: sdevalapurkar@math.harvard.edu